

Temperature for the $(2 + 1)$ -dimensional Black Hole with Non Linear Electrodynamics from the Generalized Uncertainty Principle

Alexis Larrañaga

*Universidad Nacional de Colombia. Observatorio Astronómico Nacional (OAN) and
Universidad Distrital Francisco José de Caldas. Facultad de Ingeniería. Proyecto Curricular de Ingeniería Electrónica**

Hector J. Hortua

Universidad Nacional de Colombia. Observatorio Astronómico Nacional (OAN)[†]

In this paper, we study the thermodynamical properties of the $(2 + 1)$ dimensional black hole with a non-linear electrodynamics with a negative cosmological constant, using the Generalized Uncertainty Principle (GUP). This approach shows that there is a minimum mass or remnant for the black hole, corresponding to the minimum radius of the event horizon that has a size of the order of the Planck scale. We also show that the heat capacity for this black hole is always positive.

Keywords: Black Holes Thermodynamics, Generalized Uncertainty Principle.

PACS: 04.70.Dy; 04.20.-q; 11.10.Lm

Working in $(2 + 1)$ dimensional gravity, and using as the source of the Einstein equations the stress-energy tensor of non-linear electrodynamics, Cataldo et. al. [1] found a solution with a Coulomb-like electric field (proportional to the inverse of r^2), that describes charged-AdS space when considering a negative cosmological constant.

As is well known, the thermodynamical properties of black holes are associated with the presence of the event horizon. Recently [2], it has been shown that the $(2 + 1)$ dimensional field equations for the black hole with nonlinear electrodynamics can be interpreted as the differential first law with the usual form

$$dM = TdS + \Phi dQ, \quad (1)$$

where T is the Hawking temperature that can be expressed in terms of the surface gravity at the horizon κ by

$$T = \frac{\kappa}{2\pi}. \quad (2)$$

On the other hand, in recent years the uncertainty relation that includes gravity effects, known as the Generalized Uncertainty Principle (GUP), has shown interesting results in the context of black hole evaporation [3], extending the relation between temperature and mass to scales of the order of the Planck length, $l_p = 1.61 \times 10^{-33} \text{cm}$. This treatment imply that l_p is the smallest length scale in the theory and it is related to the existence of an extreme mass (the Planck mass $m_p = 1.22 \times 10^{19} \text{GeV}$, which becomes the black hole remnant), that corresponds to the maximum possible temperature. This approach has been used recently [4] to extend the temperature-mass relation for the $(2 + 1)$ dimensional black hole with a nonlinear electric field and without cosmological constant ($\Lambda = 0$). There is also shown that there is a maximum temperature permitted for the black hole that corresponds to a horizon with size in the Planck scale.

In this paper, we investigate the thermodynamics of the $(2 + 1)$ -dimensional black hole with a nonlinear electric field reported in [1], to show how the GUP extended to gravity in spaces with cosmological constant as proposed by Arraut et. al. [3] can be used to calculate the Hawking temperature associated with the black hole. The $T(M)$ equation gives the usual relation for large masses but gets deformed when the mass becomes small. We also show how there is a minimum mass for the black hole that corresponds to a horizon with size in the Planck scale. Finally we calculate the heat capacity for this black hole, to show that it is always positive.

I. THE 3-DIMENSIONAL BLACK HOLE WITH NON-LINEAR ELECTRODYNAMICS

The metric reported by Cataldo et. al. [1] is a solution of the $(2 + 1)$ dimensional Einstein's field equations with a negative cosmological constant $\Lambda < 0$,

*Electronic address: ealarranaga@unal.edu.co

†Electronic address: hjhortuao@unal.edu.co

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi T_{\mu\nu}, \quad (3)$$

where we have used units such that $G = 1$. To obtain a Coulomb-like electric field, Cataldo et. al. used a nonlinear electrodynamics, in which the electromagnetic action does not depend only on the invariant $F = \frac{1}{4}F_{\mu\nu}F^{\mu\nu}$, but it is a function of $F^{3/4}$; and the energy-momentum tensor is restricted to be traceless. The static circularly symmetric solution obtained has the line element

$$ds^2 = -f(r) dt^2 + \frac{dr^2}{f(r)} + r^2 d\varphi^2, \quad (4)$$

where

$$f(r) = -M - \Lambda r^2 + \frac{4Q^2}{3r}. \quad (5)$$

The electric field for this solution is

$$E(r) = \frac{Q}{r^2}, \quad (6)$$

which is the standard Coulomb field for a point charge. Note that the metric depends on two parameters Q and M , that are identified as the electric charge and the mass, respectively.

A. Horizons

The horizons of this solution are defined by the condition

$$f(r) = 0 \quad (7)$$

or

$$-M - \Lambda r^2 + \frac{4Q^2}{3r} = 0. \quad (8)$$

For a cosmological constant in the range

$$-\frac{M^3}{12Q^4} \leq \Lambda < 0, \quad (9)$$

the roots of the third-order polynomial (8) are all real, and can be written as [2]

$$r_1 = -2\sqrt{-\frac{M}{3\Lambda}} \cos\left(\frac{1}{3} \cos^{-1}\left(2\frac{Q^2}{M}\sqrt{-\frac{3\Lambda}{M}}\right)\right) \quad (10)$$

$$r_2 = -2\sqrt{-\frac{M}{3\Lambda}} \cos\left(\frac{1}{3} \cos^{-1}\left(2\frac{Q^2}{M}\sqrt{-\frac{3\Lambda}{M}}\right) + \frac{2\pi}{3}\right) \quad (11)$$

$$r_3 = -2\sqrt{-\frac{M}{3\Lambda}} \cos\left(\frac{1}{3} \cos^{-1}\left(2\frac{Q^2}{M}\sqrt{-\frac{3\Lambda}{M}}\right) + \frac{4\pi}{3}\right). \quad (12)$$

Note that the form of this solution impose the condition

$$2\frac{Q^2}{M}\sqrt{-\frac{3\Lambda}{M}} \leq 1. \quad (13)$$

The equal sign defines a extreme black hole, with the maximum mass

$$M^{max} = \sqrt[3]{-12Q^4\Lambda} \quad (14)$$

and with the horizons

$$r_1 = -2\sqrt{-\frac{M^{max}}{3\Lambda}} \quad (15)$$

$$r_2 = r_3 = 2\sqrt{-\frac{M^{max}}{3\Lambda}}. \quad (16)$$

Note that the radius r_1 is negative, so it does not represent a physical horizon because for other values of the mass it is always negative. Therefore, this kind of black holes have only two physical horizons, r_2 and r_3 , that, for the extremal black hole, coincide,

$$r_2 = r_3 = r_{ext} = -\left(\frac{2Q^2}{3\Lambda}\right)^{1/3}. \quad (17)$$

In the general case, the largest radius between r_2 and r_3 corresponds to the event horizon of the black hole r_+ , while the other corresponds to the inner horizon r_- .

Using the expansion $\cos^{-1}x \approx \frac{\pi}{2} - x - \frac{1}{6}x^3$, the radii of the horizons r_2 and r_3 in equations (11) and (12) can be approximated up to the third order in

$$x = 2\frac{Q^2}{M}\sqrt{-\frac{3\Lambda}{M}}, \quad (18)$$

as

$$r_2 = r_+ \approx \sqrt{-\frac{M}{3\Lambda}} \left[\sqrt{3} - \frac{1}{3}x + \frac{\sqrt{3}}{18}x^2 + \frac{4}{81}x^3 \right] + O(x^4) \quad (19)$$

$$r_3 = r_- \approx \sqrt{-\frac{M}{3\Lambda}} \left[\frac{2}{3}x + \frac{8}{81}x^3 \right] + O(x^4). \quad (20)$$

If we consider only the first contribution in x we get

$$r_2 = r_+ \approx \sqrt{-\frac{M}{\Lambda}} - \frac{2}{3}\frac{Q^2}{M} \quad (21)$$

$$r_3 = r_- \approx \frac{4}{3}\frac{Q^2}{M}. \quad (22)$$

II. HAWKING TEMPERATURE

In this section we will deduce the Hawking temperature for this black hole by using the usual definition (2). The surface gravity can be calculated as

$$\kappa = \chi(x^\mu) a, \quad (23)$$

where a is the magnitude of the four-acceleration and χ is the red-shift factor. In order to calculate χ , we will consider a static observer, for whom the red-shift factor is just the proportionality factor between the timelike Killing vector K^μ and the four-velocity V^μ , i.e.

$$K^\mu = \chi V^\mu. \quad (24)$$

The metric (4) has the Killing vector

$$K^\mu = (1, 0, 0) \quad (25)$$

while the four-velocity is calculated as

$$V^\mu = \frac{dx^\mu}{d\tau} = \left(\frac{dt}{d\tau}, 0, 0 \right). \quad (26)$$

This gives

$$V^\mu = (f^{-1}(r), 0, 0) = \left(\frac{1}{\sqrt{-M - \Lambda r^2 + \frac{4Q^2}{3r}}}, 0, 0 \right), \quad (27)$$

and therefore, the red-shift factor is

$$\chi(r) = \sqrt{-M - \Lambda r^2 + \frac{4Q^2}{3r}}. \quad (28)$$

On the other hand, the four-acceleration is given by

$$a^\mu = \frac{dV^\mu}{d\tau}, \quad (29)$$

that has components

$$a^0 = a^\varphi = 0 \quad (30)$$

$$a^r = -\Lambda r - \frac{2}{3} \frac{Q^2}{r^2}. \quad (31)$$

and therefore, the magnitude of the four-acceleration is

$$a = \sqrt{g_{\mu\nu} a^\mu a^\nu} = \frac{-\Lambda r - \frac{2}{3} \frac{Q^2}{r^2}}{\sqrt{-M - \Lambda r^2 + \frac{4Q^2}{3r}}}. \quad (32)$$

Then, the surface gravity at the event horizon is given by the absolute value

$$\kappa_+ = \left| -\Lambda r - \frac{2}{3} \frac{Q^2}{r^2} \right|_{r=r_+}. \quad (33)$$

Note that for $r = r_{ext}$, the surface gravity is $\kappa(r_{max}) = 0$, i.e. that the extreme black hole has zero Hawking temperature,

$$T(M^{max}) = 0. \quad (34)$$

Using the approximate forms for r_+ and r_- given by equations (19) and (20), we can write the Hawking temperature as a function of the mass. For the event horizon we have, up to second order in x ,

$$T_+(M) = \frac{\kappa_+}{2\pi} = \frac{1}{2\pi} \left| -\Lambda r_+ - \frac{2}{3} \frac{Q^2}{r_+^2} \right| \quad (35)$$

$$T_+(M) \approx \frac{1}{2\pi} \left| -\Lambda \sqrt{-\frac{M}{3\Lambda}} \left[\sqrt{3} - \frac{1}{3}x \right] - \frac{2}{3} Q^2 \left(-\frac{M}{3\Lambda} \right)^{-1} \left[\sqrt{3} - \frac{1}{3}x \right]^{-2} \right| \quad (36)$$

$$T_+(M) \approx \frac{1}{2\pi} \left| \sqrt{-M\Lambda} - \sqrt{-\frac{M\Lambda}{3}} \frac{1}{3}x + \frac{2\Lambda Q^2}{M} \left(3 - \frac{2x}{\sqrt{3}} \right)^{-1} \right| \quad (37)$$

$$T_+(M) \approx \frac{1}{2\pi} \left| \sqrt{-M\Lambda} - \sqrt{-\frac{M\Lambda}{3}} \frac{1}{3}x + \frac{2\sqrt{3}\Lambda Q^2}{M(3\sqrt{3} - 2x)} \right| \quad (38)$$

$$T_+(M) \approx \frac{1}{2\pi} \left| \sqrt{-M\Lambda} - \frac{2}{3} \frac{\Lambda Q^2}{M} + \frac{2\sqrt{-M\Lambda} Q^2}{3\sqrt{-\frac{M^3}{\Lambda} - 4Q^2}} \right|. \quad (39)$$

Note that, for $Q = 0$, this expression becomes

$$T_+^{BTZ}(M) = \frac{\sqrt{-M\Lambda}}{2\pi}, \quad (40)$$

that corresponds to the temperature at the horizon of the static non-charged BTZ black hole [5]. On the other hand, for the inner horizon r_- we have

$$T_-(M) = \frac{\kappa_-}{2\pi} = \frac{1}{2\pi} \left| -\Lambda r_- - \frac{2}{3} \frac{Q^2}{r_-^2} \right| \quad (41)$$

$$T_-(M) \approx \frac{1}{2\pi} \left| -\Lambda \sqrt{-\frac{M}{3\Lambda}} \frac{2}{3}x - \frac{2}{3} Q^2 \left[\sqrt{-\frac{M}{3\Lambda}} \frac{2}{3}x \right]^{-2} \right| \quad (42)$$

$$T_-(M) \approx \frac{1}{2\pi} \left| -\Lambda \sqrt{-\frac{M}{3\Lambda}} \frac{2}{3}x + \frac{9\Lambda Q^2}{2Mx^2} \right| \quad (43)$$

$$T_-(M) \approx \frac{1}{2\pi} \left| -\frac{4}{3} \frac{\Lambda Q^2}{M} - \frac{3M^2}{8Q^2} \right|. \quad (44)$$

III. HAWKING RADIATION AND THE GENERALIZED UNCERTAINTY PRINCIPLE

Now we will consider the GUP and the Hawking radiation derived from it. We will show how the GUP will produce a deformation in the temperature-mass relation when considered close to the Planck lenght. As stated by Arraut et. al. [3] the GUP can be stated, in the case of a non-zero cosmological constant and in units with $c = \hbar = 1$, by the relation

$$\Delta x \Delta p \gtrsim \frac{1}{2} + \frac{G}{2} (\Delta p)^2 - \frac{\gamma}{3\Lambda} \frac{1}{(\Delta p)^2}, \quad (45)$$

where G is the gravitational constant, Λ is cosmological constant and γ is a constant factor which accounts for the fact that the GUP equation is dealing with orders of magnitudes estimates. When comparing the GUP results with the above Hawking temperature, we will see that γ is expected to be of the order of 1. Since the Planck length l_p can be written as

$$l_p^2 = G\hbar = \frac{\hbar^2}{m_p^2}, \quad (46)$$

where m_p is the mass of Planck, the GUP can be written as

$$\Delta x \Delta p \gtrsim \frac{1}{2} + \frac{l_p^2}{2} (\Delta p)^2 - \frac{\gamma}{3\Lambda} \frac{1}{(\Delta p)^2}, \quad (47)$$

or

$$\Delta x \Delta p \gtrsim \frac{1}{2} + \frac{1}{2} \frac{(\Delta p)^2}{m_p^2} - \frac{\gamma}{3\Lambda} \frac{1}{(\Delta p)^2}. \quad (48)$$

To apply this uncertainty principle to the black hole evaporation process consist in identifying the Δx with the event horizon radius r_+ and the momentum Δp with the Hawking temperature up to a 2π factor [?] ($\Delta P \sim 2\pi T$). Therefore, we can write the GUP as a fourth order equation for the temperature,

$$r_+ 2\pi T = \frac{1}{2} + \frac{2\pi^2 T^2}{m_p^2} - \frac{\gamma}{3\Lambda} \frac{1}{4\pi^2 T^2} \quad (49)$$

For high temperatures, the GUP equation can be approximated by

$$r_+ 2\pi T = \frac{1}{2} + \frac{2\pi^2 T^2}{m_p^2} \quad (50)$$

that corresponds to a quadratic polynomial for T ,

$$\frac{4\pi^2}{m_p^2} T^2 - 4\pi r_+ T + 1 = 0 \quad (51)$$

From which it follows that the temperature-mass relation is

$$T(M) = \frac{m_p^2 r_+}{2\pi} \left[1 \pm \sqrt{r_+^2 - \frac{1}{m_p^2}} \right]. \quad (52)$$

Note that the argument in the square root defines a minimum radius for the event horizon,

$$r_+^{min} = \frac{1}{m_p} = l_p, \quad (53)$$

i.e. that r_+ can not be smaller than the Planck length. This length corresponds to a minimum mass or remnant for the black hole of the order of

$$M^{min} = \Lambda l_p^2. \quad (54)$$

Using equation (39), the emperature of the black hole corresponding to this mass is

$$T_+ (M^{min}) \approx \frac{1}{2\pi} \left| |\Lambda| l_p - \frac{2}{3} \frac{Q^2}{l_p^2} + \frac{2 |\Lambda| l_p Q^2}{3 |\Lambda| l_p^3 - 4 Q^2} \right|. \quad (55)$$

Note that this expression gives, in the $Q = 0$ case, a minimum mass for the BTZ black hole,

$$T_+^{BTZ} (M^{min}) = \frac{|\Lambda| l_p}{2\pi}. \quad (56)$$

Finally, the heat capacity for the $\Lambda = 0$ black hole can be calculated using equation (52),

$$C_Q = \left(\frac{\partial M}{\partial T} \right)_Q \quad (57)$$

$$= \frac{1}{2\pi m_p^2} \frac{\sqrt{r_+^2 - \frac{1}{m_p^2}}}{2r_+^2 - \frac{1}{m_p^2}} \left(-\frac{1}{2\Lambda} \sqrt{-\frac{\Lambda}{M}} + \frac{2}{3} \frac{Q^2}{M^2} \right)^{-1}. \quad (58)$$

Note that the heat capacity of this black hole is negative, $\frac{\partial T}{\partial M} < 0$, when

$$r_+^2 < \frac{1}{2m_p^2} = \frac{l_p^2}{2}, \quad (59)$$

but condition (53) impose that the minimum radius is $r_+^{min} = l_p$. Therefore the heat capacity of this black hole is always positive.

IV. CONCLUSION

We have studied the thermodynamics of the $(2+1)$ dimensional black hole with non-linear electrodynamics and with a negative cosmological constant using the Generalized Uncertainty Principle. This gives a minimum mass or remnant for the black hole, that corresponds to a maximum Hawking temperature depending only on the electric charge Q . The solution with the maximum temperature is obtained when the black hole has a size of the order of the Planck scale (minimum horizon).

Equation (52) gives the temperature-mass relation, and as is shown, it gives the standard Hawking temperature (39) for small masses, but gets deformed for masses close to M^{min} . Finally, the heat capacity of this black hole is always positive. This analysis confirms that Planck lenght seems to be the smallest lenght in nature, even in $(2+1)$ dimensions.

-
- [1] M. Cataldo, N. Cruz, S. del Campo and A. Garcia. *Phys. Lett. B* **484**, 154 (2000)
 - [2] A. Larranaga and L. A. Garcia. arXiv: 0811.3368 [gr-qc]
 - [3] I. Arraut, D. Batic and M. Nowakowski. arXiv: 0810.5156 [gr-qc]
 - [4] A. Larranaga. arXiv: 0811.3368 [gr-qc]
 - [5] M. Akbar, Chin. Phys. Lett. **24** (5) (2007) 1158